

ENTROPIC NONEXTENSIVITY: A POSSIBLE MEASURE OF COMPLEXITY

Constantino Tsallis

Centro Brasileiro de Pesquisas Físicas, Xavier Sigaud 150, 22290-180 Rio de Janeiro-RJ, Brazil
Department of Physics, University of North Texas, P.O. Box 311427, Denton, Texas 76203, USA
Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, New Mexico 87501, USA
tsallis@cbpf.br

**"Beauty is truth, truth beauty," – that is all
 Ye know on earth, and all ye need to know.**

(John Keats, May 1819)

An updated review [1] of nonextensive statistical mechanics and thermodynamics is colloquially presented. Quite naturally the possibility emerges for using the value of $q - 1$ (entropic nonextensivity) as a simple and efficient manner to provide, at least for some classes of systems, some characterization of the degree of what is currently referred to as complexity [2]. A few historical digressions are included as well.

Keywords: Nonextensive statistical mechanics; Entropy; Complexity; Multifractals; Power laws.

I. INTRODUCTION

Would you say that Newtonian mechanics is *universal*? We almost hear a big and unanimous *yes* coming from practically all nineteenth-century physicists. And yet, they would all be wrong! Indeed, we know today that if the involved masses are very small, say that of the electron, Newtonian mechanics has to be replaced by quantum mechanics. And if the involved speeds are very high, close to that of light in vacuum, special relativity has to be used instead. And if the masses are very large, like that of the Sun or of a galaxy, general relativity enters into the game. It is our present understanding that only in the $(\hbar, 1/c, G) \rightarrow (0, 0, 0)$ limit, Newtonian mechanics appears to be strictly correct. Now, what about Boltzmann-Gibbs statistical mechanics and standard thermodynamics? Are they *universal*? An attempt to answer such question and to clarify its implications constitutes the main goal of the present effort. Let us see what Ludwig Boltzmann has to tell us about this. In the first page of the second part of his *Vorlesungen über Gastheorie*, he qualifies the concept of ideal gas by writing: "When the distance at which two gas molecules interact with each other noticeably is vanishingly small relative to the average distance between a molecule and its nearest neighbor – or, as one can also say, when the space occupied by the molecules (or their spheres of action) is negligible compared to the space filled by the gas – ...". Were Boltzmann our contemporary, he would have perhaps told us that he was addressing systems involving what we nowadays call *short range* interactions! Several decades ago, Laszlo Tisza [3], in his *Generalized Ther-*

modynamics, writes "The situation is different for the additivity postulate [...], the validity of which cannot be inferred from general principles. We have to require that the interaction energy between thermodynamic systems be negligible. This assumption is closely related to the homogeneity postulate [...]. From the molecular point of view, additivity and homogeneity can be expected to be reasonable approximations for systems containing many particles, provided that the intramolecular forces have a short range character", when referring to the usual thermodynamic functions such as internal energy, entropy, free energy, and others. Also, Peter T. Landsberg [4], in his *Statistical Mechanics and Thermodynamics*, writes "The presence of long-range forces causes important amendments to thermodynamics, some of which are not fully investigated as yet.". More, during a visit to Rio de Janeiro some years ago, he told me that *such restrictions should be in the first pages of all books on thermodynamics... but they are not!* The title of the 1993 Nature article [5] by John Maddox focusing on black holes is "When entropy does not seem extensive". What is the viewpoint on such matters of our colleagues mathematicians? Distinguished expert in nonlinear dynamical systems, Floris Takens writes [6] "The values of p_i are determined by the following dogma: if the energy of the system in the i^{th} state is E_i and if the temperature of the system is T then: $p_i = \exp\{-E_i/kT\}/Z(T)$, where $Z(T) = \sum_i \exp\{-E_i/kT\}$ [...] This choice of p_i is called Gibbs distribution. We shall give no justification for this dogma; even a physicist like Ruelle disposes of this question as "deep and incompletely clarified". It is known that mathematicians sometimes use the word *dogma* when they do not have the theorem! As early as 1964, Michael E. Fisher, and later on with David Ruelle as well as with Joel L. Lebowitz, addressed in detail such questions [7]. They established, for instance, that a d -dimensional classical system (say a fluid) including two-body interactions that are nonsingular at the origin and decay, at long distances, like $r^{-\alpha}$, exhibits standard thermostistical behavior if $\alpha/d > 1$. An interesting question remains open: what happens when $\alpha/d \leq 1$? This is the case of all self-gravitating systems (e.g., galaxies, black holes), a fact which explains well known difficulties in using standard thermodynamics to address them. These difficulties are not without relation with the case of a single hydrogen atom ($d = 3$ and $\alpha = 1$),

for which no tractable Boltzmann-Gibbs thermostatical calculations are possible (diverging partition function) unless we confine the hydrogen atom in a box [8]. Of course, no such difficulties are encountered if we focus on a single harmonic oscillator, or on a single spin in an external magnetic field, or on a Lennard-Jones fluid ($d = 3$ and $\alpha = 6$), or even on a neutral plasma, where the Coulombian interaction is "dressed" thus becoming a thermostatically innocuous, exponentially decaying interaction. No severe anomalies appear in such cases, all relevant sums and integrals being finite and perfectly defined. More generally speaking, a variety of recent analytical and numerical results (see [9] and references therein) in classical systems (with no singularities at the potential origin or elsewhere) suggest that, at the appropriately defined ultimate $t \rightarrow \infty$ thermal equilibrium, the internal energy *per particle* grows, when the number of particles N grows, like $\tilde{N} \equiv [N^{1-\alpha/d} - \alpha/d]/[1 - \alpha/d]$. We easily verify that $\tilde{N}$ approaches a finite constant if $\alpha/d > 1$, diverges logarithmically if $\alpha/d = 1$, and diverges like $N^{1-\alpha/d}$ if $0 \leq \alpha/d < 1$, thus recovering, for $\alpha = 0$, the well known Molecular Field Approximation scaling with N (usually englobed within the coupling constant whenever dealing with magnetic models such as the Ising ones, among many others). We trivially see that the standard thermodynamical *extensivity* is lost in such cases. The physical behavior for the marginal case $\alpha/d = 1$ is intimately related to the mathematical fact that some relevant sums that are *absolutely convergent* on the extensive side and *divergent* on the nonextensive one, become *conditionally convergent* in that case. This enables us to understand why the amount of calories to be provided to a table in order to increase its temperature in one degree *only* depends on its weight and material (iron, wood), whereas the amount of Coulombs we must provide to a capacitor to generate a one Volt potential difference at its ends *also* depends on its *shape*! Indeed, the relevant interaction in the latter being the permanent dipole-dipole one (hence $d = \alpha = 3$), the capacity depends on whether the capacitor is a slab, round, cylindrical or otherwise. Such behavior dramatically contrasts with the extensivity usually focused on in textbooks of thermodynamics. In fact, it is after all quite natural that self-gravitating systems provided the first physical application of the formalism we are addressing here. Indeed, physicist Angel Plastino and his son astronomer Angel R. Plastino showed in 1993 [10] that by sufficiently departing from BG thermostatics, it is possible to overcome an old difficulty, namely to have the physically desirable feature that total energy, entropy and mass be *simultaneously finite*.

Let us now try to deepen our quest for understanding the restrictions within which Boltzmann-Gibbs statistical mechanics is the appropriate formalism for describing a (fantastic!) variety of systems in nature. Are long-range interactions the only mechanism which creates anomalies? The answer seems to be *no*. Indeed, several indica-

tions exist which suggest that mesoscopic dissipative systems (like granular matter) also are thermostatically anomalous: the velocity distribution measured in a variety of computational experiments is, as argued by Leo P. Kadanoff et al [11], Y.-H. Taguchi and H. Takayasu [12], Hans J. Herrmann [13] and others, far from being Maxwell's Gaussian. On general grounds, multifracally structured systems also tend to exhibit anomalies. And I believe that it would be surprise for very few that the same happened with systems based on strongly non-markovian microscopic memory (e.g., a memory function decreasing towards the past as a slow enough power-law).

As an attempt to theoretically handle at least some of these anomalies, I proposed in 1988 [14] the generalization of the Boltzmann-Gibbs formalism by postulating a nonextensive entropy S_q (See Sections I and II) which recovers the usual logarithmic measure as the $q = 1$ particular case. Before going further, let us describe S_q , its properties and historical context.

II. THE NONEXTENSIVE ENTROPY S_q

A. Properties

In the case of a discrete number W of microscopic configurations, S_q is given by [14]

$$S_q \equiv k \frac{1 - \sum_{i=1}^W p_i^q}{q - 1} \quad \left(\sum_{i=1}^W p_i = 1; k > 0 \right) \quad (1)$$

(similar expressions correspond to the cases where we have a continuum of microscopic configurations, or when the system is a quantum one).

In the limit $q \rightarrow 1$ we have $p_i^{q-1} = e^{(q-1) \ln p_i} \sim 1 + (q-1) \ln p_i$, hence $S_1 = -k \sum_{i=1}^W p_i \ln p_i$, which is the usual Boltzmann-Gibbs-Shannon expression. Also, it can be shown that

(i) $S_q \geq 0$, and, for $q > 0$, equals zero in the case of certainty (i.e., when all probabilities vanish but one which equals unity);

(ii) S_q attains its extremum (maximum for $q > 0$ and minimum for $q < 0$) for equiprobability (i.e., $p_i = 1/W$), thus becoming $S_q = k[W^{1-q} - 1]/[1 - q]$. This expression becomes, for $q = 1$, $S_1 = k \ln W$, which is the formula graved on Boltzmann's tombstone in the Central Cemetery in Vienna;

(iii) S_q is concave (convex) for $q > 0$ ($q < 0$), which is the basis for thermodynamic stability;

(iv) If two systems A and B are independent in the sense of the theory of probabilities (i.e., $p_{ij}^{A+B} = p_i^A \times p_j^B$), then

$$\frac{S_q(A+B)}{k} = \frac{S_q(A)}{k} + \frac{S_q(B)}{k} + (1-q) \frac{S_q(A)}{k} \frac{S_q(B)}{k}, \quad (2)$$

hence, superextensivity, extensivity or subextensivity occurs when $q < 1$, $q = 1$ or $q > 1$ respectively;

(v) If the W states of a system are divided into W_L and W_M states ($W_L + W_M = W$), and we define $p_L = \sum_{W_L \text{ states}} p_i$ and $p_M = \sum_{W_M \text{ states}} p_m$ ($p_L + p_M = 1$), then

$$S_q(\{p_i\}) = S_q(p_L, p_M) + p_L^q S_q(\{p_i/p_L\}) + p_M^q S_q(\{p_m/p_M\}) \quad (3)$$

which generalizes Shannon's celebrated formula;

(vi) It can be easily checked that $-d(\sum_i p_i^x)/dx|_{x=1} = -\sum_i p_i \ln p_i$. Sumiyoshi Abe, in Funabashi, made in 1997 [15] a simple but very deep remark. He considered the so-called *Jackson derivative*. The mathematician F. Jackson generalized, one century ago, the usual differential operator. He defined $D_q f(x) \equiv [f(qx) - f(x)]/[qx - x]$, which reproduces d/dx in the limit $q \rightarrow 1$. Abe's observation consists in the following easily verifiable property, namely $-D_q(\sum_i p_i^x)/dx|_{x=1} = S_q$! Since it is clear that Jackson derivative "tests" how the function $f(x)$ "reacts" under dilatation of x in very much the same way the standard derivative "tests" it under translation of x , Abe's remark opens a wide door to physical insight onto thermodynamics. It was so perceived, I believe, by Murray Gell-Mann, of the Santa Fe Institute. Indeed, one year ago, during a conference in Italy, he asked me whether I would go to conjecture that systems with relevant symmetries *other* than say translation or dilatation invariances would need entropies *other* than S_1 or S_q . My answer was *probably yes*. Indeed, there are symmetries that appear in general relativity or in string theory that are characterized by a sensible amount of parameters. The concept of thermodynamic information upon such systems, i.e., the appropriate entropy could well be a form involving essentially a similar number of parameters. If so, then symmetry would control entropy, and S_q would only be the beginning of a presumably long and fascinating story! Since we have learnt from Gell-Mann and others, how deeply symmetry controls energy, we could then say that symmetry controls thermodynamics, that science which nothing is but a delicate balance between energy and entropy. More precisely, symmetry would then determine (see also [16]) the *specific microscopic form* that Clausius thermodynamic entropy would take for *specific* classes of systems. Symmetry controlling thermodynamics! If such a conjectural scenario turned out to be correct, this would not be to displease Plato with his unified view of truth and beauty!

(vii) Along the lines of Claude E. Shannon's theorem, it has been shown by Roberto J. V. dos Santos, in Macao, Brazil [17], that a set of conditions generalizing (through properties (iv) and (v)) that imposed by Shannon uniquely determines S_q . Analogously, Abe recently showed [18] that consistently generalizing Khinchin's set of conditions, once again a unique solution emerges, namely S_q .

B. Brief review of the labyrinthic history of the entropies

The entropy and the entropic forms have haunted physicists, chemists, mathematicians, engineers and others since long! At least since Clausius coined the word and wrote "*The energy of the universe is constant, its entropy tends to increase*" (See Appendix). The whole story is a labyrinth plenty of rediscoveries. After Ludwig Boltzmann and Josiah Willard Gibbs introduced and first analyzed, more than one century ago, their respective expressions [20] of the entropy in terms of microscopic quantities, many generalizations and reformulations have been proposed and used. Let us mention some of the most relevant ones. John von Neumann introduced a quantum expression for the entropy in terms of the density operator; when the operator is diagonalized, the traditional expression (herein referred, because of the functional form, to as the Boltzmann-Gibbs entropy, although these two scientists used to work with different, though consistent, expressions [20]) is recovered. Shannon reinvented, more than half a century ago, a binary version of the B-G functional form and interpreted it in terms of information and communication theory. It is no doubt very interesting to notice that in *The mathematical theory of communication* he wrote, in reference to the specific form he was using, that "*This theorem, and the assumptions required for its proof, are in no way necessary for the present theory. It is given chiefly to lend a certain plausibility to some of our later definitions. The real justification of these definitions, however, will reside in their implications.*". In 1957, Edwin Jaynes attributed to the whole formalism a very generic, and still controversial in spite of an enormous amount of practical applications, anthropomorphic shape. Since those years, people working in cybernetics, information theory, complexity and nonlinear dynamical systems, among others, have introduced close to 20 different entropic forms! This number has almost doubled, this time mainly because of physicists, since the 1988 paper [14] appeared.

In 1960, A. Renyi proposed [21] a form which recovers that of Shannon as a special case. His form is *always extensive*, but *not always concave (or convex)* with regard to the probability distributions. It seems that it was a rediscovery; indeed, according to I. Csizsar's 1974 critical survey [22], that form had essentially already been introduced by Paul-Marcel Schutzenberger in 1954 [23]. In 1967, presumably for cybernetic goals, J. Harvda and F. Charvat introduced [24], although with a different prefactor (fitted for binary variables), the entropic form herein noted S_q , and studied some of its mathematical properties. One year later, I. Vajda further studied [25] this form, quoting Harvda and Charvat. In 1970, in the context of information and control, Z. Daroczy rediscovered [26] this form (he quotes neither Harvda and Charvat, nor Vajda). Later on, in 1975, B.D. Sharma and D.P. Mittal introduced [27] and studied some mathematical proper-

ties of a two-parameter form which recovers *both* Renyi's and S_q as particular instances! In his 1978 paper in Reviews of Modern Physics, A. Wehrl mentions [28] the form of S_q , quoting Daroczy, but not Harvda, Charvat, Vajda, Sharma and Mittal. In 1988, in the Physics community, a new rediscovery, by myself this time. Indeed, inspired by multifractals, I postulated [14] the form S_q as a possible new path for generalizing Boltzmann-Gibbs statistical mechanics; through optimization of S_q I obtained an equilibrium-like distribution which generically is a *power-law*, and reproduces Boltzmann's celebrated factor as the particular limit $q \rightarrow 1$. In my 1988 paper, I quoted only Renyi's entropy, having never taken notice of all these generalized forms, whose existence (and that of many others!) I have been discovering along the years. In fact, Renyi's entropy can be shown [14] to be a monotonically increasing function of S_q . However, concavity is not preserved through monotonicity. So what?, since the optimizing distribution will be exactly the same. Well, it happens that statistical mechanics is much more than an equilibrium-like distribution optimizing an entropy under some generic constraints! It includes a variety of other relevant points, such as the role played by fluctuations, linear responses, thermodynamical stability, to name but a few. Nobody would, I believe, try to reformulate BG statistical mechanics using, instead of the BG entropy, say its *cube*, just because it is a monotonically increasing function of the BG one! Let us also mention that, within various scientific and technological communities, close to 30 (!) different entropic forms have been considered along the last half a century.

Let us summarize some of what we have said. The BG entropy, S_1 , is both *concave* and *extensive*, the Renyi entropy, S_q^R , is *extensive* but *nonconcave* (for all $q > 0$; concave only for $0 < q \leq 1$), the one focused on herein, S_q , is *nonextensive* but *concave* (for all $q > 0$), and the Sharma-Mittal entropy is *nonextensive* and *nonconcave*. The latter is determined by *two* parameters and contains the other three as particular cases. The Renyi entropy and S_q are determined by *one* parameter (q) and share a common case, namely the BG entropy, which is nonparametric. Since Renyi's, Sharma-Mittal's and the present entropy depend from the probability set through one and the same expression, namely $\sum_i p_i^q$, they are all three related through simple functions. In particular,

$$S_q^R \equiv k \frac{\ln \sum_i p_i^q}{1-q} = k \frac{\ln[1 + (1-q)S_q/k]}{1-q} \quad (\forall q), \quad (4)$$

with $S_1^R = S_1 = -k \sum_i p_i \ln p_i$.

For completeness, let us close this section by mentioning that expressions of the type $\sum_i p_i^q$ have also been, long ago, discussed on mathematical grounds by Hardy, Littlewood and Polya [19] (and, for $q = 2$, by the Pythagoreans!).

III. EQUILIBRIUM-LIKE STATISTICS

The next step is to follow along Gibbs path. For instance, to formally obtain the "equilibrium" distribution associated with the canonical ensemble (i.e., a conservative system at "equilibrium" with a thermostat, where the physical meaning of this state will turn out to be, as we shall see later on, close to a stationary one), we need to impose a constraint on total energy. We shall adopt [29] that $\langle \mathcal{H} \rangle_q \equiv \sum_i P_i \epsilon_i = U_q$, where U_q is fixed and finite, $\{\epsilon_i\}$ is the set of energy levels of the system Hamiltonian $\mathcal{H}$, and the *escort* distribution $\{P_i\}$ is defined as $P_i \equiv p_i^q / [\sum_j p_j^q]$; from now on we refer to $\langle \dots \rangle_q$ as the *normalized q -expectation value*. At the present stage, a plethora of mathematical reasons exist for adopting this particular generalization of the familiar concept of *internal energy*. Not the least of these reasons is the fact that such definition enables relevant sums and integrals to be *finite*, which would otherwise diverge (e.g., the second moment of Lévy distributions diverges, whereas the second q -moment with q appropriately chosen is *finite*). We can say that in this sense the theory becomes *regularized*. The optimization of S_q with this constraint (to which we associate a Lagrange parameter β) leads to

$$p_i = [1 - (1-q)\beta_q(\epsilon_i - U_q)]^{1/(1-q)} / Z_q \quad (\beta_q \equiv \beta / Z_q^{1-q}) \quad (5)$$

with $Z_q \equiv \sum_j [1 - (1-q)\beta_q(\epsilon_j - U_q)]^{1/(1-q)}$ (See Fig. 1). We easily verify that $q \rightarrow 1$ recovers the celebrated, *exponential* Boltzmann factor

$$p_i = e^{-\beta \epsilon_i} / Z_1 \quad (Z_1 \equiv \sum_j e^{-\beta \epsilon_j}). \quad (6)$$

For $q > 1$ a *power-law* tail emerges; for $q < 1$ the formalism imposes a high-energy *cutoff*, i.e., $p_i = 0$ whenever the argument of the power function becomes negative. One comment is worthy: this distribution is generically *not* an exponential law, i.e, it is generically *not* factorizable (under sum in the argument), and nevertheless is *invariant* under the choice of zero energy for the energy spectrum! (this is in fact one of the aside benefits of defining the constraints in terms of *normalized* distributions like the escort ones).

Eq. (5) can be rewritten in the following convenient form:

$$p_i = [1 - (1-q)\beta'_q \epsilon_i]^{1/(1-q)} / Z'_q \quad (7)$$

with $Z'_q \equiv \sum_j [1 - (1-q)\beta'_q \epsilon_j]^{1/(1-q)}$, where β' is a simple function of β_q and U_q .

It can be shown [29], along the lines of the 1991 paper with Evaldo M.F. Curado [29], that this formalism satisfies, for *arbitrary* q , $1/T = \partial S_q / \partial U_q$ ($T \equiv 1/(k\beta)$), and the entire Legendre transform structure of standard thermodynamics. Moreover, the relations which provide

U_q and $F_q = U_q - TS_q$ in terms of Z_q (i.e., the connection between the microscopic and macroscopic descriptions of the system), remain basically the *same* as the usual ones, the role played by the logarithmic function $\ln x$ being now played by $(x^{1-q} - 1)/(1 - q)$. Finally, various important thermostistical theorems can be shown to be q -invariant. Among them, we must mention the Boltzmann H-theorem (macroscopic time irreversibility, i.e., essentially the second principle of thermodynamics, first tackled within this context by Ananias M. Mariz in Natal, Brazil [30]), Ehrenfest theorem (correspondence principle between classical and quantum mechanics), Einstein 1910 factorizability of the likelihood function in terms of the entropy (this q -generalization was first realized in 1993 during a private discussion with Manuel O. Caceres, from Bariloche, Argentina), Onsager reciprocity theorem (microscopic time reversibility), Kramers and Wannier relations (causality), and, apparently, even Pesin relation for nonlinear dynamical systems, although the last one remains at the level of plausibility (no proof available). The q -generalization of the fluctuation-dissipation theorem was first addressed by Anna Chame and Evandro V.L. de Mello, in Niteroi, Brazil [31], and then, along with the generalization of Kubo's linear response theory, by A.K. Rajagopal, in the Naval Research Laboratory in Washington [32]. Since then, a variety of important statistical mechanical approaches such as the Bogolyubov inequality, the many-body Green function and path integral formalisms have been generalized by Renio S. Mendes in Maringa, Brazil, Ervin K. Lenzi in Rio de Janeiro, and collaborators [32]. Nonextensive versions of Fermi-Dirac and Bose-Einstein statistics have been discussed [33] by Fevzi Buyukkilic, Dogan Demirhan and Ugur Tirnakli in Izmir, Turkey, and their collaborators. Finally, it is certainly important to mention that, very recently, Rajagopal and Abe have done an exhaustive review [34] of apparently *all* the traditional methods for obtaining the BG equilibrium distribution, namely the Balian-Balazs counting method within a microcanonical basis, the Darwin-Fowler steepest descent method, and the Khinchin's large-numbers method. They q -generalized all three, and systematically obtained the same power-law distribution obtained [14,29] above through the Gibbs' method, i.e., the optimization of an appropriately constrained entropy.

In order to see the nonextensive formalism at work, let us address [35] an important question which long remained without satisfactory solution, namely, why Lévy distributions are ubiquitous in nature, in a manner which by all means is similar to the ubiquity of Gaussians? To introduce the point, let us first address the Gaussian case. If we optimize the BG entropy S_1 by imposing a fixed and *finite* average $\langle x^2 \rangle$, we obtain a Gaussian one-jump distribution. By convoluting N such distributions we obtain a N -jump distribution which also is a Gaussian. From this immediately follows that $\langle x^2 \rangle(N) \propto N$. If we consider that N is proportional to time, we have the central result of Einstein's 1905 celebrated paper on

Brownian motion, which provided at the time strong evidence in favor of Boltzmann's ideas. We may reword this result by saying that the foundation of Gaussians in nature lies upon two pillars, namely the BG entropy and the central limit theorem. What about Lévy distributions? The difficulty comes from the fact that all Lévy distributions have *infinite* second moment, due to their long tail ($\propto 1/|x|^{1+\gamma}$ with $0 < \gamma < 2$). Consequently, what simple auxiliary constraint to use along with the optimization of the entropy? In 1994, Pablo Alemany and Damian H. Zanette in Bariloche, Argentina, settled [36] the basis for the solution, namely optimization of S_q while imposing a *finite* q -expectation value of x^2 . The one-jump distribution thus obtained is proportional to $[1 - (1 - q)\beta x^2]^{1/(1-q)}$, which respectively reproduces the Gaussian, Lorentzian (*Cauchy* for the mathematicians) and completely flat distributions for $q = 1, 2$ and 3 . For $q < 1$ the support is compact, and for $q > 1$ a power-law tail is obtained. The proof was completed one year later [35], when it was argued that the Lévy-Gnedenko central limit theorem was applicable for $q > 5/3$, thus approaching, for $N \gg 1$ precisely Lévy distributions for the N -jump distribution. We may then summarize this result by saying that the foundation of Lévy distributions in nature also has two pillars, namely S_q and the Lévy-Gnedenko theorem. It follows that $\gamma = 2$ for $q \leq 5/3$ and $\gamma = (3 - q)/(q - 1)$ for $5/3 < q < 3$. Since γ can also be interpreted as the fractal dimension to be associated with Lévy walks, this turned out to be the first *exact* connection between nonextensive statistical mechanics and scaling. Similar ideas have been successfully put forward by Plastino and Plastino [37] and by Lisa Borland [38] and others, concerning nonlinear as well as fractional-derivative Fokker-Planck-like equations. In all these various types of anomalous diffusion, the Gaussian solution associated with Jean Baptiste Joseph Fourier's bicentennial heat equation is recovered as the $q = 1$ limit. Let me also mention that Hermann Haken, in Stuttgart, and collaborators have recently applied related ideas within a theoretical model of the human brain [39].

In the same vein which led to the above discussion of Lévy distributions, if we optimize S_q imposing, besides normalization of $p(x)$, finite values for the normalized q -expectation values of *both* x and x^2 , we obtain

$$p_q(x) \propto [1 - (1 - q)(\beta^{(1)}x + \beta^{(2)}x^2)]^{1/(1-q)}, \quad (8)$$

where $\beta^{(1)}$ and $\beta^{(2)}$ are determined by the constraints. This distribution contains, as particular cases, a considerable amount of well (and not so well) known distributions, such as the exponential, the Gaussian, the Cauchy-Lorentz, the Edgeworth [40] (for $\beta^{(1)} = 1, \beta^{(2)} = 0$, and $x \geq 0$), the r - and the Student's s - ones [41], among others. Let us end with a short and amusing historical remark concerning that which can be considered as the most famous of all nontrivial distributions, namely the normal one. In contrast with what almost everybody would naively think, it was [42] first introduced by Abraham De Moivre in 1733, then by Pierre Simon de Laplace

in 1774, then by Robert Adrain in 1808, and finally by Carl Friedrich Gauss in 1809, nothing less than 76 years after its first publication!

IV. MICROSCOPIC DYNAMICAL FOUNDATION

To have a deeper understanding of what this generalized formalism means, it is highly convenient to focus on the mixing properties.

A. Mixing in one-dimensional dissipative maps

Let us now address a very simple type of system, which will nevertheless provide important hints about the physical and mathematical significance of the present formalism. Let us consider the well known logistic map, namely

$$x_{t+1} = 1 - ax_t^2 \quad (t = 0, 1, 2, \dots; 0 \leq a \leq 2) \quad (9)$$

For $a < a_c = 1.40115519\dots$, the attractors are *finite* cycles (fixed points, cycle-2, cycle-4, etc) and the Lyapunov exponent (hereafter noted λ_1) is almost everywhere *negative* (i.e., *strongly insensitive* to the initial conditions and any intermediate rounding). For $a_c \leq a \leq 2$, the attractor is chaotic for most values (i.e., it has an *infinite* number of elements, analogously to an irrational number) and the λ_1 exponent is consistently *positive* (i.e., *strongly sensitive* to the initial conditions and any intermediate rounding). There is however, in the interval $[0, 2]$, an infinite number of values of a for which $\lambda_1 = 0$. What happens with the sensitivity to the initial conditions in those marginal cases? Let us focus on this problem by first recalling the definition of λ_1 . If two initial conditions x_0 and x'_0 differ by the small amount Δx_0 , we can follow the time evolution of Δx_t through the quantity $\xi \equiv \lim_{\Delta x_0 \rightarrow 0} (\Delta x_t / \Delta x_0)$. The most frequent situation is that $\xi = e^{\lambda_1 t}$, which defines the Lyapunov exponent λ_1 . We can trivially check that $\xi(t)$ is the solution of $d\xi/dt = \lambda_1 \xi$. We have conjectured [43] that, whenever λ_1 *vanishes* for this map (and similar ones), the controlling equation becomes $d\xi/dt = \lambda_q \xi^q$, hence $\xi = [1 + (1 - q)\lambda_q t]^{1/(1-q)}$. This result recovers the usual one at the $q = 1$ limit; also, it defines a *generalized* Lyapunov coefficient λ_q , which, like λ_1 , inversely scales with time, but now within a law which generically is a *power*, instead of the standard *exponential*. The two paradigmatic cases are $q < 1$ (with $\lambda_q > 0$), hereafter referred to as *weakly sensitive* to the initial conditions, and $q > 1$ (with $\lambda_q < 0$), hereafter referred to as *weakly insensitive* to the initial conditions. Within this unified scenario, we would have *strong chaos* for $q = 1$ and $\lambda_1 > 0$ (i.e., *exponential mixing*) and *weak chaos* for $q < 1$, $\lambda_1 = 0$ and $\lambda_q > 0$ (i.e., *power-law mixing*). All these various possibilities have indeed been observed for the logistic map. If we start with $x_0 = 0$, we have (i) $\lambda_1 < 0$ for $a = a_c - 10^{-3}$, (ii) $\lambda_1 > 0$ for $a = a_c + 10^{-3}$, (iii)

$\lambda_1 = 0$, $q > 1$ and $\lambda_q < 0$ for a doubling-period bifurcation (e.g., $a = 3/4$) as well as for a tangent bifurcation (e.g., $a = 7/4$), and finally (iv) $\lambda_1 = 0$, $q < 1$ and $\lambda_q > 0$ at the edge of chaos (i.e., $a = a_c$). The last case is, for the present purposes, the most interesting one. This power-law mixing was first observed and analyzed by P. Grassberger, A. Politi, H. Mori and collaborators [44]. Its importance seems to come mainly from the fact that, in the view of many authors, "life appeared at the edge of chaos"! Moreover, this type of situation seems to be a paradigm for vast classes of the so-called *complex systems*, much in vogue nowadays. Let us say more about this interesting point. The precise power-law function ξ indicated above appears to be the *upper bound* of a complex time-dependence of the sensitivity to the initial conditions, which includes considerable and ever lasting fluctuations. From a log-log representation we can numerically get the slope and, since this is to be identified with $1/(1 - q)$, we obtain the corresponding value q^* . For the logistic map we have obtained [43] $q^* = 0.24\dots$. If we apply these concepts to a more general map, like $x_{t+1} = 1 - a|x_t|^z$ ($0 \leq a \leq 2$; $z > 1$), we can check, and this is long known, that the edge of chaos depends on z . More precisely, when z increases from 1 to infinity, a_c increases from 1 to 2. If for a_c we check the sensitivity to the initial conditions, we obtain that q^* increases from $-\infty$ to close to 1 (though smaller than 1), when z increases from 1 to infinity. Simultaneously, the fractal dimension d_f of the chaotic attractor increases from 0 to close to 1 (though smaller than 1). If instead of the logistic family of maps, we use some specific family of circle maps also characterized by an inflexion exponent z , it can be shown that, once again, q^* increases with z for all studied values of z ($z \geq 3$). The interesting feature in this case is that $d_f = 1$ for *all* these values of z . What we learn from this is that the index q characterizes essentially the *speed* of the mixing, and only indirectly how "filled" is the phase space within which this mixing is occurring.

Let us now address, for maps like the logistic one, a *second* manner of obtaining q^* , this time based on the multifractal geometry of the chaotic attractor. This geometry can be, and usually is, characterized by the so called $f(\alpha)$ function, a down-ward parabola-like curve which generically is below (and tangential to) the bissector in the (α, f) space, is concave, its maximal value is d_f , and, in most of the cases, vanishes at two points, namely α_{min} and α_{max} (typically, $0 < \alpha_{min} < d_f < \alpha_{max}$). For a fractal, one expects, $\alpha_{min} = d_f = \alpha_{max} = f(\alpha_{min}) = f(d_f) = f(\alpha_{max})$. For a so-called *fat* (multi)fractal, d_f attains the value of the Euclidean space within which the (multi)fractal is embedded (this is the case of the family of circle maps mentioned above). Marcelo L. Lyra, in Maceio, and myself used (imitating the successful Ben Widom style!) in 1998 [43] some simple scaling arguments and obtained the following relation:

$$\frac{1}{1-q^*} = \frac{1}{\alpha_{min}} - \frac{1}{\alpha_{max}} \quad (q^* < 1) \quad (10)$$

This is a kind of fascinating relation since it connects the power-law sensitivity of nonlinear dynamical systems with purely geometrical quantities. It has been numerically checked for at least three different one-dimensional dissipative maps. Its analysis for two- or more-dimensional systems, as well as for conservative ones, would of course be very enlightening and remains to be done (as do quantum analogs of this problem). Furthermore, it does not contain in a generic way the $q = 1$ limit. Indeed, it seems that, for its validity, it is implicitly assumed that $f(\alpha_{min}) = f(\alpha_{max}) = 0$. A more general expression could be something like $1/(1-q^*) = 1/[\alpha_{min} - f(\alpha_{min})] - 1/[\alpha_{max} - f(\alpha_{max})]$; the generic $q = 1$ limit could then correspond to $\alpha_{min} = f(\alpha_{min})$. The second manner we referred above goes as follows. We can numerically construct the $f(\alpha)$ function associated with the chaotic attractor, measure α_{min} and α_{max} , and, through the above connection, calculate q^* . This procedure yields, for the standard logistic map, $q^* = 0.2445\dots$, thus reproducing the number we previously obtained from the sensitivity to the initial conditions. The higher precision we indicate here comes from the fact that, for the logistic-like family maps, it is long known that $\alpha_{max} = z \alpha_{min} = \frac{\ln 2}{\ln \alpha_F}$, where α_F denotes the corresponding Feigenbaum constant. This constant being known with many digits for $z = 2$, we can infer q^* with quite high precision.

Let us finally address now a *third* manner of obtaining q^* , this time directly from the nonextensive entropy S_q . It is known that, for large classes of nonlinear dynamical systems, the rate of loss of information as time increases can be characterized by the so called Kolmogorov-Sinai entropy (hereafter noted K_1). This quantity is defined, strictly speaking, through a *single-trajectory* procedure, based on the frequencies of appearance, in increasingly long strips, of symbolic sequences associated with partitions of the phase space. However, in apparently very large (if not all) classes of problems, this single-trajectory procedure can be replaced by an *ensemble-based* one, whose numerical implementation is by far simpler computationally than the implementation of algorithms based on sequences. We shall restrict ourselves to this ensemble-based procedure. Let us illustrate, on the logistic map, the steps to be followed in order to calculate K_1 . We choose a value of a for which λ_1 is positive, say $a = 2$ (whose corresponding Lyapunov exponent is $\lambda_1 = \ln 2$). We then partition the accessible phase space ($-1 \leq x \leq 1$, in our example) in $W \gg 1$ little windows, and choose (in any way), inside one of those windows, N initial values for x_0 . We then apply the map onto each of those initial conditions, which, as time evolves, will start spreading within the accessible phase space. At time t , we will observe a set $\{N_i(t)\}$ of points inside the W windows ($\sum_{i=1}^W N_i(t) = N$, $\forall t$). We next define a set of probabilities $p_i \equiv N_i/N$ ($i = 1, 2, \dots, W$), which en-

able in turn the calculation of $S_1(t) = -\sum_{i=1}^W p_i \ln p_i$. This entropy starts, by construction, from zero at $t = 0$, and tends to linearly increase as time goes on, finally saturating at a value which, of course, cannot exceed $\ln W$. The Kolmogorov-Sinai-like entropy rate is then defined through $K_1 \equiv \lim_{t \rightarrow \infty} \lim_{W \rightarrow \infty} \lim_{N \rightarrow \infty} S_1(t)/t$. We expect, according to the Pesin equality, $K_1 = \lambda_1$. Indeed, we numerically obtain $K_1 = \ln 2$ (see Fig. 2(a)). We can analogously define a generalized quantity, namely $K_q \equiv \lim_{t \rightarrow \infty} \lim_{W \rightarrow \infty} \lim_{N \rightarrow \infty} S_q(t)/t$. And we can numerically verify that, for say $a = 2$, K_q vanishes (diverges) for any value of $q > 1$ ($q < 1$), *being finite only for $q = 1$* . Let us now turn onto the edge of chaos, i.e., $a = a_c$. Vito Latora and Andrea Rapisarda, in Catania, Italy, Michel Baranger, at MIT, and myself [43], have numerically verified the same structure, excepting for the fact that now K_q vanishes (diverges) for any value $q > q^*$ ($q < q^*$), *being finite only for $q = q^* = 0.24\dots$* ! (see Fig. 2(b)). Tirnakli et al [45] have also verified the validity of this structure for various typical values of z within the logistic-like family of maps.

Let us summarize this section by emphasizing that we have seen that it is possible to arrive to one and the same nontrivial value of q^* through *three* different and suggestive roads, namely sensitivity to the initial conditions, multifractal structure, and rate of loss of information. By so doing, we obtain insight onto the mathematical and physical meaning of q and its associated nonextensive formalism. Moreover, a pleasant unification is obtained with the concepts already known for standard, exponential mixing. A very interesting question remains open though, namely the generalization of Pesin equality. It might well be that, as conjectured in [43], K_q equals λ_q (or, more generically, an appropriate average of $\lambda_q(x)$, whenever nonuniformity is present). However, a mathematical basis for properties such as this one is heavily needed. Some recent numerical results along these lines by Paolo Grigolini, in Denton, Texas, and collaborators [46] seem promising. In the same vein, it is known that whenever *escape* exists, as time evolves, out of a specific region of the phase space of a chaotic system, K_1 becomes larger than λ_1 , the difference being precisely the *escape rate* [47]. All these quantities are defined through *exponential* time-dependences. What happens if all three are *power* laws instead? A natural speculation emerges, namely that perhaps a similar relation exists for $q \neq 1$.

B. Mixing in many-body dissipative maps

In the previous section we described a variety of interesting nonextensive connections occurring in one-dimensional dissipative systems. A natural subsequent question then is what happens when the dissipative system has many, say $N \gg 1$, degrees of freedom? Such is the case of essentially all the models exhibiting Per Bak's self-organized criticality (SOC). Francisco Tamarit

and Sergio Cannas, in Cordoba, Argentina, and myself have discussed [48] this issue for the Bak-Sneppen model [49] for biological evolution. By numerically studying the time evolution of the Hamming distance for increasingly large systems, a power-law was exhibited. By identifying the exponent with $1/(1 - q^*)$, the value $q^* \simeq -2.1$ was obtained. Totally analogous results are obtained for the Suzuki-Kaneko model for imitation games (e.g., bird singings) [50]. It would be of appreciable importance if the same numbers were achieved through the other two procedures, namely the determination of the $f(\alpha)$ function and of the loss of information. Such consistency would be of great help for further understanding. These tasks remain, however, to be done.

C. Mixing in many-body Hamiltonian systems

Let us now address what might be considered as the *Sancta Sanctorum* of statistical mechanics, namely the systems that Boltzmann himself was mainly focusing on, i.e., classical many-body Hamiltonian systems. We know that no severe thermostistical anomalies arrive for N -body d -dimensional systems whose (say two-body) interactions are neither singular at the origin nor long-ranged. A most important example which violates *both* restrictions is of course Newtonian gravitation. We shall here restrict to a simpler case, namely interactions that are well behaved at the origin, but which can be long-ranged. For specificity, let us assume a two-body potential which (attractively) decays at long distances like $1/r^\alpha$ with $0 \leq \alpha$ (for historical reasons we use α for this exponent; however, it has clearly nothing to do with the α appearing in the characterization of multifractals previously discussed). The limit $\alpha \rightarrow \infty$ corresponds to very short interactions; the other extreme ($\alpha = 0$) corresponds to a typical Mean Field Approximation. No thermostistical basic difficulties are expected for $\alpha > \alpha_c$ but the situation is quite different for $0 \leq \alpha \leq \alpha_c$. As already mentioned, Fisher and collaborators [7] established that $\alpha_c = d$ for classical systems ($\alpha_c \geq d$ is expected for quantum systems). Therefore, the nontrivial case is $0 \leq \alpha/d \leq 1$. Two physically different situations are to be understood, namely what happens at the $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty}$ situation, and what happens at the $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty}$ one. The two orderings are expected to be essentially equivalent for $\alpha/d > 1$, and dramatically different otherwise. Let us first focus on the $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty}$ ordering. This should correspond to the traditional, BG-like, concept of thermal equilibrium, though with anomalous scalings. Let us be more specific and consider a fluid thermodynamically described by

$$G(T, p, N) = U(T, p, N) - TS(T, p, N) + pV(T, p, N), \quad (11)$$

hence, using the quantity $\tilde{N}$ introduced in Section I,

$$\frac{G(T, p, N)}{N\tilde{N}} = \frac{U(T, p, N)}{N\tilde{N}} - \frac{T}{\tilde{N}} \frac{S(T, p, N)}{N} + \frac{p}{\tilde{N}} \frac{V(T, p, N)}{N}. \quad (12)$$

In the thermodynamic limit $N \rightarrow \infty$ we expect to obtain *finite* quantities as follows:

$$g(\tilde{T}, \tilde{p}) = u(\tilde{T}, \tilde{p}) - \tilde{T}s(\tilde{T}, \tilde{p}) + \tilde{p}v(\tilde{T}, \tilde{p}), \quad (13)$$

with $\tilde{T} \equiv T/\tilde{N}$ and $\tilde{p} \equiv p/\tilde{N}$. So, when $\alpha/d > 1$, $\tilde{N}$ is a constant, and we recover the usual concepts that G , U , S and V are *extensive* variables since they scale like N , whereas T and p are *intensive* variables since they are invariant with respect to N . If, however, $0 \leq \alpha/d \leq 1$, the extensivity-intensivity concepts lose their usual simple meanings. Indeed, there are now *three*, and *not two*, thermodynamical categories, namely the *energetic* ones (G , F , U , ...) which scale like $N\tilde{N}$, the *pseudo-intensive* ones (T , p , H , ...) which scale like $\tilde{N}$, and their conjugate *pseudo-extensive* variables (S , V , M , ...) which scale like N (*if and only if* expressed in terms of the pseudo-intensive variables). I can unfortunately not prove the general validity of the above scheme, but it has been shown to be true in *all* the models that have been numerically or analytically studied [51] (one- and two-dimensional Ising and Potts magnets, one-dimensional bond percolation and XY coupled rotators, one-, two- and three-dimensional Lennard-Jones-like fluids, among others). Also, for the case $\alpha = 0$, we recover the traditional scalings of Mean Field Approximations, where the Hamiltonian is *artificially* made extensive by dividing the coupling constant by N . By so doing, we are obliged to pay a conceptually high price, namely to have microscopic couplings which depend on N ! This long standing tradition in magnetism has, for very good reasons, never been adopted in astronomy: we are not aware of any astronomer dividing the universal gravitational constant G by any power of N in order to artificially make extensive a Hamiltonian which clearly is not! The thermodynamical scheme that has been described above escapes from such unpleasant criticism. I sometimes refer to the above discussion of long-ranged systems as *weak violation* of BG statistical mechanics. Indeed, it becomes necessary to conveniently rescale the thermodynamic variables (using $\tilde{N}$) but the $q = 1$ approach remains the adequate one, i.e., energy distributions still are of the exponential type.

Let us now address the much more complex $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty}$ situation. At the present moment, the reliable informations that we have are mainly of numerical nature, more precisely computational molecular dynamics. A paradigmatic system has been intensively addressed during last years, namely a chain of coupled inertial planar rotators (ferromagnetic XY-like interaction), the interaction decaying as $1/r^\alpha$. The system being classical, it is thermodynamically extensive for $\alpha > 1$ and nonextensive for $0 \leq \alpha \leq 1$. Important informations were established by Ruffo, in Florence, and collaborators [52], who focused on the $\alpha = 0$ case. If we note E_N the

total energy associated with N rotators, it can be shown that a second order phase transition occurs at the critical point $e \equiv E_N/(N\tilde{N}) = 3/4 \equiv e_c$. For $e > e_c$ the rotators are nearly decoupled and for $e < e_c$ they tend to clusterize. Not surprising, e_c depends on α , and vanishes above a critical value of α which is larger than unity. For $\alpha > 1$, nothing very anomalous occurs. But, in the interval $0 \leq \alpha \leq 1$, curious phenomena happen on *both* sides of $e_c(\alpha)$.

For e above e_c (say $e = 5$), the (conveniently scaled) maximum Lyapunov exponent $\lambda_{max}(N)$ goes to zero as $N^{-\kappa}$, where $\kappa(\alpha)$ was numerically established, by Celia Anteneodo and myself [53] (see Fig. 3). Nothing like this is observed for $\alpha > 1$, where the numerical evidence strongly points that $\lim_{N \rightarrow \infty} \lambda_{max}(N) > 0$. This of course suggests that, for a thermodynamically large system, mixing is not exponential, but possibly a power-law, i.e., weak mixing, as discussed above for the one-dimensional maps. The $\kappa(\alpha)$ dependence has been recently studied by Andrea Giansanti, in Rome, and collaborators for $d = 2$ and 3 as well. If plotted against α/d (see Fig.3), an universal curve seems to emerge. This evidence is reinforced by the fact that it is possible to analytically prove, for this and other models, that $\kappa(0) = 1/3$. It is consistently obtained $\kappa = 0$ for $\alpha/d \geq 1$.

Let us now focus on the anomalies *below* e_c . The case $e = 0.69$ for $\alpha = 0$ has been and is being further studied in detail. The maximal Lyapunov exponent is here positive in the limit $N \rightarrow \infty$ (for all $0 < e < e_c$ in fact) *but*, suggestively enough, the usual (BG) canonical ensemble equilibrium is preceded by a long, metastable-like, *quasi stationary state* (QSS), whose duration τ diverges with N ! This means that, if N is of the order of the Avogadro number, during a time that might be longer than the age of the universe, only this curious QSS will be observable. So, we better understand it! As can be seen in Fig. 4, the average kinetic energy per particle e_k depends on both N and time t . Only in the $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty}$ the BG canonical temperature T is attained. If we take the limits the other way around a completely different result is obtained, namely e_k approaches a finite value below T , and τ diverges like N^ρ . It becomes possible to speculate that $\rho(\alpha)$ decreases from a finite positive value to zero, when α/d increases from zero to unity; thereafter, we of course expect $\rho = 0$, the system being then a well behaved BG one. To make this scenario more robust, consistent evidence is available concerning diffusion of the $\alpha = 0$ rotators. The numerical study of this phenomenon has shown [52,56] an average second moment of the angle increasing like t^β , where β depends on both N and t . For fixed N , $\beta > 1$ (i.e., superdiffusion) for a long time until, at $t = \tau_D$ (D stands for *diffusion*), makes a crossover to unity (normal diffusion). Analogously to what was described above, τ_D increases like N^σ . Once again, it is possible to speculate that $\sigma(\alpha)$ decreases from some finite positive value to zero when α/d increases from zero to unity. The door remains open to whether there is a simple connection between σ and ρ . All these re-

sults enable to conjecture that a scenario close to what is shown in Fig. 5 could indeed occur for some physical systems. Before closing this section, let me emphasize that what Fig. 4 strongly suggests is the *possible* validity of the zeroth principle of thermodynamics even *out* of a canonical BG scenario! Indeed, for large enough N , systems can share the same "temperature", thus being at thermal equilibrium, though this equilibrium is *not* the familiar one. This is quite remarkable if we take into account that there is nowadays quite strong numerical evidence that the distribution of velocities in such systems is far from Gaussian, i.e., the familiar Maxwellian distribution of velocities is by all means violated.

V. COMPARISON WITH EXPERIMENTAL RESULTS

During this last decade many types of comparisons have been done between the present theory and experimental data. They have different epistemological status, and range from simple fittings (with sometimes clear, sometimes rather unclear physical insight), up to completely closed theories, with no fitting parameters at all. Let me review a few among them.

A. Turbulence

X.-P. Huang and C.F. Driscoll exhibited in 1994 [57] some quite interesting nonneutral electronic plasma experiments done in a metallic cylinder in the presence of an axial magnetic field. They observed a turbulent axisymmetric metaequilibrium state, the electronic density radial distribution of which was measured. Its average (over typically 100 shots) monotonically decreased with the radial distance, disappearing at some radius sensibly *smaller* than the radius of the container (i.e., a *cut-off* was observed). They also proposed four phenomenological theories trying to reproduce the experimentally observed profile. The *Restricted Minimum Enstrophy* one provided a quite satisfactory first-approximation fitting. Bruce M. Boghosian, in Boston, showed in 1996 [58] that the Huang and Driscoll successful attempt *precisely* corresponds to the optimization of S_q with $q = 1/2$, the necessary cut-off naturally emerging from the formalism. Since then, improved versions of his calculation [14,59], as well as controversial arguments [60], have been published. The merit of first connecting the present formalism to turbulence remains however doubtless.

In recent months, Fernando M. Ramos and collaborators, in São José dos Campos, Brazil [61], Toshihiko Arimitsu and N. Arimitsu, in Tokyo [62], as well as Christian Beck, in London [63], have exhibited very interesting connections with fully developed turbulence. Beck's theory is a closed one, with no fitting parameters. He

obtained a quite impressive agreement with the experimental distribution of the differences (at distance r) of radial velocities, including its well known slight skewness (see Fig. 6). The entropic parameter is given by $1/(q-1) = 1 + \log_2(r/\eta)$ where η is the viscosity. The fact that q is not universal certainly reminds us the criticality of the two-dimensional XY ferromagnet, with which nonextensivity presents in fact various analogies.

B. High-energy collisions

Ignacio Bediaga and collaborators [64], in Rio de Janeiro, have worked out, along Rolf Hagedorn's 1965 lines [65], a phenomenological theory for the distribution of transverse momenta in the hadronic jets emerging from electron-positron annihilation after central collisions at energies ranging from 14 to 161 GeV. Indeed, since early ideas of Fermi, and later of Field and Feynman, a thermodynamical equilibrium scenario has been advanced for this distribution of transverse momenta. Hagedorn developed a full theory based on the BG distribution. The success was only partial. Indeed, a central ingredient of the physical picture is that higher collision energies *do not increase the transverse momenta temperature T* but increase instead the number of involved bosons that are produced (like water boiling at higher flux of energy, where only the rate of vapor production is increased, but not the temperature). It is this central physical expectation that was not fulfilled by Hagedorn's calculation; indeed, to fit the curves associated with increasingly high energies, increasingly high values for T become necessary. Furthermore, an *inflexion point* clearly emerges in the distribution, which by no means can be reproduced by the BG exponential distribution. The adequation of only two parameters (T and q) have enabled Bediaga et al [64] to fit amazingly large sets of experimental data (see Fig. 7). And, in order to make the expected picture complete, the central demand of T being independent from the collision energy is fulfilled!

Grzegorz Wilk, in Warsaw, and collaborators [66], as well as D.B. Walton and J. Rafelski in Tucson [67], have provided further evidences of the applicability of the present ideas to high energy physics, where nonmarkovian processes and long-range interactions are commonly accepted hypothesis. Consistent evidence has also been provided by D.B. Ion and M.L.D. Ion, in Bucharest, very especially in hadronic scattering [68].

C. Solar neutrino problem

Piero Quarati and collaborators, in Torino and Cagliari, have been advancing since 1996 [69] an interesting possibility concerning the famous solar neutrino problem. Indeed, it is known that calculations within the so-called Standard Solar Model (SSM) provide a neutrino

flux to be detected on the surface of the Earth which is roughly the *double* of what is actually measured in several laboratories around the world. This enigmatic discrepancy intrigues the specialists since over 20-30 years ago. At least two, *non exclusive*, possible causes are under current analysis. One of them concerns the nature of neutrinos: they could oscillate in such a way that only part of them would be detectable by the present experimental devices. The second one addresses the possibility that the SSM could need to be sensibly improved in what concerns the production of neutrinos at the solar core plasma. It is along this line that Quarati's suggestion progresses. Indeed, the neutrino flux is related to the total area of the so-called Gamow peak, which is in turn due to the product of the thermal equilibrium BG distribution function (which decreases with energy) and the penetration factor (which increases with energy). The position of the peak is at energies 10 times larger than kT , therefore only the far *tail* of the thermal distribution is concerned. Quarati et al argue that very slight departures from $q = 1$ (of the order of 0.01) are enough to substantially modify (close to the desired factor two) the area of the Gamow peak. They also show that this degree of departure remains within the experimental accuracy of other independent measurements, such as the helioseismographic ones. This slight nonextensivity would be related to quite plausible nonmarkovian processes and other anomalies.

D. Others

Many other phenomena have been, during recent years, tentatively connected to the present nonextensivity. Let me briefly mention some of them. The experimental, clearly non Gaussian, distribution of velocities in various systems has been shown to be very satisfactorily fitted by $q \neq 1$ curves. Such is the case of the distribution (observed from the COBE satellite) of peculiar velocities of spiral galaxies, shown [70], in 1998 by Quarati, myself and collaborators, to be very well fitted by $q \simeq 0.24$. Such is also the case of the distribution that James Glazier, in Notre Dame University, and collaborators [71], measured for the horizontal velocities of *Hydra Vulgaris* living in physiological solution. They were remarkably well fitted using $q \simeq 1.5$. Such is finally the case of the computationally simulated granular matter involving mesoscopic inelastic collisions. Indeed, the distribution function that Y.-H. Taguchi and H. Takayasu used to fit their 1995 data [12] precisely is a typical $q > 1$ one.

In 1995 [72], F.C. Sa Barreto, in Belo Horizonte, Brazil, E.D. Loh, in Michigan State University, and myself advanced the possibility of explaining very tiny departures from the black-body radiation Planck's law for fitting the COBE satellite data for the cosmic background microwave radiation. Indeed, values of q departing from unity by the order of 10^{-5} were advanced. This possibility has since then be further analyzed by a variety

of scientists, the conclusion still remaining basically the same. Only more precise (perhaps 10 times more precise would be enough) experimental data would confirm or reject such a possibility, physically motivated by present or ancient subtle effects of gravitation, which could even lead ultimately to a modification of our understanding of the nature of spacetime (at very small scales, it could very well be noncontinuous and even nondifferentiable, in contrast with our usual perceptions!).

Molecules like CO and O_2 in some hemoproteins can be dissociated from their natural positions by light flashes, as many classical experiments have shown. Under specific circumstances, these molecules then tend to re-associate with a time-dependent rate. The experimental rate has been shown, in 1999 by G. Bemski, R.S. Mendes and myself [73], to be very well fitted by the solutions of a nonextensive-inspired differential equation. The intrinsic fractality of such proteins is believed to be the physical motivation for such approach. Numbers of citations of scientific publications can also be well fitted with $q > 1$ curves, consistently with S. Denisov's 1997 recovering, within the present formalism, of the so-called Zipf-Mandelbrot law for linguistics [74]. Oscar Sotolongo-Costa, in La Habana, and collaborators [75], as well as K.K. Gudima, in Caen, France, and collaborators [76], have recently invoked the present formalism for describing multifragmentation.

Last but not least, it is worthy mentioning that the present framework has induced [77] quite quicker versions of the Simulated Annealing methods for global optimization. This methodology has found successful applications in theoretical chemistry, in particular for studying proteins, their folding and related phenomena. Such procedures are currently being used by John E. Straub, at the Boston University, and collaborators, by Kleber C. Mundim, in Salvador, Brazil, Donald E. Ellis, in Chicago, and collaborators, by Yuko Okamoto, in Okazaki, Ulrich H.E. Hansmann, in Houghton, and collaborators, among others [78].

VI. FINAL COMMENTS

Many issues remain, at the present moment, partially or fully open to better understanding concerning the present attempt of adequately generalizing Boltzmann-Gibbs statistical mechanics and thermodynamics. These include (i) the direct checking of the energy distribution which generalizes the BG factor (and consistently the connection between q and (α, d) for long-range interacting Hamiltonian systems (both finite and large), and similar connections for nonconservative ones); (ii) appropriate understanding within the renormalization group framework (Alberto Robledo, in Mexico, has recently initiated this line [79]; see also [80]); (iii) possible firm connections of this formalism with the quantum group formalism for generalizing quantum mechanics (the great

analogies that these two nonextensive formalisms exhibit are since long being explored, but not yet deeply understood); (iv) possible connections with the Lévy-like problematic reviewed recently [81] by George M. Zaslavsky; (v) the strict conditions (in terms of the $N \rightarrow \infty$ and $t \rightarrow \infty$ limits) under which the zeroth-principle of thermodynamics (i.e., the criterium for thermal equilibrium) holds; (vi) the microscopic and mathematical interpretation of the direct and escort distributions (which possibly reflect the choice of a (multi)fractal description, or of a description in terms of Lebesgue-integrable quantities); (vii) the general connection between symmetry and entropic form adapted to measure information about systems having that particular symmetry; (viii) the rigorous basis for the generalization herein described of concepts such as Kolmogorov-Sinai entropy, Lyapunov exponents, Pesin equality, escape rates, etc; (ix) the possibility of having, for *long-range* interacting systems, a special value for q which, through the use of appropriately rescaled variables, would enable a new kind of extensivity in the sense that $S_q(A + B) = S_q(A) + S_q(B)$, where the *large* subsystems A and B would of course be *not independent* (in the sense of theory of probabilities) but, on the contrary, strongly coupled; (x) the connection between the index q appearing in properties such as the sensitivity to the initial conditions, and that appearing in properties such as the microscopic distribution of energies (the most typical values for the former appear to be *smaller* than unity, whereas they are *larger* than unity for the latter); (xi) the direct checking of the type of mixing which occurs in the phase-space of many-body long-range interacting large systems; (xii) the deep connection with quantum entanglement through its intrinsic nonlocality. The last point certainly appears as very promising and relevant. An important step forward was very recently done by Abe and Rajagopal [82]; indeed, they succeeded in recovering, through use of the present formalism, the $x = 1/3$ Peres (necessary and sufficient) condition, which is known to be the *strongest* one (stronger than Bell inequality) for having local realism (some degree of separability of the full density matrix into those associated with subsystems A and B) for the bipartite spin- $\frac{1}{2}$ system (Werner-Popescu state). The probability of this approach being fundamentally correct is enhanced nowadays by contributions produced from quite different viewpoints [83].

As we can see, the number of important issues still needing enlightening is quite large. This makes the enterprise but more stimulating! However, caution is recommended in spite of the sensible amount of objective successes of the proposal. Indeed, given the enormous impact of the thermodynamical concepts in Physics, one should address all these points with circumspection. Nevertheless, an important new perspective seems to be already acquired. Nikolai S. Krylov showed, half a century ago, that the key concept in the foundation of standard statistical mechanics is not ergodicity but mixing. Indeed, he showed that the Lyapunov exponents essen-

tially control the relaxation times towards BG equilibrium. The entropy associated with such systems is vastly known to be the BG logarithmic one. What emerges now is an even more fascinating possibility, namely the *type* of mixing seems to determine the microscopic *form* of the entropy to be used. If the mixing is of the exponential type (strong chaos), then the entropy of course is the BG one (i.e., $q = 1$). If the mixing is of the power type (weak chaos), then the mixing exponent ($1/(1 - q)$ apparently) would determine the anomalous value of q to be used for the entropy, which would in turn determine the thermal equilibrium distribution, as well as all of its thermodynamical consequences! The kingdom of the exponentials would then be occasionally replaced by the kingdom of the power laws, with their relevance in biology, economics and other complex systems!

I thank warm hospitality at the University of North Texas, at the Santa Fe Institute and at MIT, where this paper was written. I am especially grateful to M. Gell-Mann, who made my visit to the Santa Fe Institute possible, stimulating and pleasant. My sincere thanks go also to L. E. Reichl whose editorial interest and dedication greatly improved the entire Proceedings volume, and very especially the present manuscript. Work in my group at the Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, is supported in part by PRONEX/FINEP, CNPq and FAPERJ (Brazilian Agencies).

APPENDIX

Scene at the restaurant

AU LABYRINTHE DES ENTROPIES

– Would you have some fresh entropies today, for me and my friends?

– Absolutely Sir ! We have them extensive or not, with definite concavity or not, nonnegative defined or otherwise, quantum, classical, relative, cross or mutual, included in several others with a small supplement, composable or not, expansible or not, totally optimized or a little rare, even completely out of equilibrium... single-trajectory-based or ensemble-based...

You can have them at the good old Boltzmann magnificent style, dorée à la Gibbs, very subtle, or von Neumann..., with pepper à la Jaynes, or the popular Shannon, oh, my God, I was forgetting the esoteric, superb, macroscopic Clausius, the surprising Fisher, the refined Kolmogorov-Sinai, à la Kullback-Leibler for comparison, Renyi with multifractal dressing, with cybernetic sauce à la Harvda and Charvat, Vajda and Daroczy, or even the all-taste Sharma and Mittal...

In 1988 we started serving them with Brazilian touch, if you wish to try, it leaves a tropical arrière-gout in your mouth! And since then, our chefs have introduced not less than ten new recipes... Curado, with exponentials, Anteneodo with tango flavor, Plastino's, excellent as family dish, Landsberg, Papa, Johal with curry, Borges and

Roditi, Rajagopal and Abe...

They are all delicious ! How many do I serve you today?

FIGURES

Fig. 1 - Equilibrium-like probability versus energy for typical values of q , namely, from top to bottom at low energies, $q = 0, 1/4, 1/2, 2/3, 1, 3, \infty$ (the latter collapses onto the ordinate at the origin, and vanishes for all positive energies). For $q = 1$ we have Boltzmann exponential factor; for $q > 1$ we have a power-law tail; for $q < 1$ there is a cutoff above which the probability vanishes.

Fig. 2 - Time evolution of S_q : (a) $a = 2$ ($q^* = 1$); (b) $a = a_c$ ($q^* = 0.2445$; R measures the degree of nonlinearity of the curves in the relevant intermediate region).

Fig. 3 - Coupled planar rotators: κ versus α/d ($d = 1$: [53]; $d = 2, 3$: [54]). The solid line is a guide to the eye.

Fig. 4 - Coupled planar rotators ($\alpha = 0, d = 1$): Time evolution of T . From Latora and Rapisarda [55]).

Fig. 5 - Conjectural time dependence of the probability p of having an energy E for classical systems ($\tilde{N} \equiv N^* + 1$).


Fig. 6 - Distribution of radial velocity differences in fully developed turbulence (from top to bottom: $r/\eta = 3.3, 23.6, 100$). From Beck [63].


Fig. 7 - Distribution of hadronic transverse momenta: (a) cross-section (dotted line: Hagedorn's theory; solid lines: $q \neq 1$ theory); (b) Fitting parameters q and T_0 . From Bediaga et al [64].


-
- [1] The present review corresponds to the inaugural talk delivered at the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000). The Proceedings are edited by P. Grigolini, C. Tsallis and B.J. West, and will appear in *Chaos, Fractals and Solitons* (2001).
- [2] M. Gell-Mann, *The quark and the jaguar* (W.H. Freeman, New York, 1994).
- [3] L. Tisza, *Annals Phys.* **13**, 1 (1961) [or in *Generalized thermodynamics*, (MIT Press, Cambridge, 1966), p. 123].
- [4] P.T. Landsberg, *Thermodynamics and Statistical Mechanics*, (Oxford University Press, Oxford, 1978; also Dover, 1990), page 102.
- [5] J. Maddox, *Nature* **365**, 103 (1993).
- [6] F. Takens, *Structures in dynamics - Finite dimensional deterministic studies*, eds. H.W. Broer, F. Dumortier, S.J. van Strien and F. Takens (North-Holland, Amsterdam, 1991), page 253.
- [7] M.E. Fisher, *Arch. Rat. Mech. Anal.* **17**, 377 (1964); *J. Chem. Phys.* **42**, 3852 (1965); *J. Math. Phys.* **6**, 1643 (1965); M.E. Fisher and D. Ruelle, *J. Math. Phys.* **7**, 260 (1966); M.E. Fisher and J.L. Lebowitz, *Commun. Math. Phys.* **19**, 251 (1970).
- [8] R.H. Fowler, *Statistical Mechanics* (Cambridge University, Cambridge, 1936), Chap. XIV.
- [9] C. Tsallis, in *Nonextensive Statistical Mechanics and Thermodynamics*, Series *Lecture Notes in Physics*, eds. S. Abe and Y. Okamoto (Springer-Verlag, Heidelberg, 2000), in press; M. Antoni, S. Ruffo and A. Torcini, in the Proceedings of the Conference "The Chaotic Universe" (Rome-Pescara, February 1999) [cond-mat/9908336].
- [10] A.R. Plastino and A. Plastino, *Phys. Lett. A* **174**, 384 (1993).
- [11] Yunson Du, Hao Li and L. P. Kadanoff, *Phys. Rev. Lett.* **74**, 1268 (1995).
- [12] Y.-H. Taguchi and H. Takayasu, *Europhys. Lett.* **30**, 499 (1995).
- [13] H.J. Herrmann, private communication.
- [14] C. Tsallis, *J. Stat. Phys.* **52**, 479 (1988). See <http://tsallis.cat.cbpf.br/biblio.htm> for further bibliography on the subject. A set of mini-reviews is contained in "Nonextensive Statistical Mechanics and Thermodynamics", eds. S.R.A. Salinas and C. Tsallis, *Braz. J. Phys.* **29** (1999) [http://sbf.if.usp.br/WWW_pages/Journals/BJP/Vol29/Num1/index.htm].
- [15] S. Abe, *Phys. Lett. A* **224**, 326 (1997).
- [16] L.E. Reichl, *A Modern Course in Statistical Physics* (Wiley Interscience, New York, 2nd. edition, 1998).
- [17] R.J.V. dos Santos, *J. Math. Phys.* **38**, 4104 (1997).
- [18] S. Abe, *Phys. Lett. A* **271**, 74 (2000).
- [19] G. Hardy, J.E. Littlewood and G. Polya, *Inequalities* (Cambridge University Press, Cambridge, 1952).
- [20] J.L. Lebowitz, *Physics Today* **46**, 32 (1993) and *Physica A* **263**, 516 (1999).
- [21] A. Renyi, in *Proc. Fourth Berkeley Symposium*, 1960, Vol. 1 (University of California Press, Berkeley, Los Angeles, 1961), 547; *Probability Theory* (North-Holland, 1970) and references therein.
- [22] I. Csiszar, *Information measures: A critical survey*, in *Transactions of the Seventh Prague Conference on Information Theory, Statistical Decision Functions, Random Processes and the European Meeting of Statisticians*, 1974 (Reidel, Dordrecht, 1978) page 73.
- [23] M.P. Schutzenberger, *Contribution aux applications statistiques de la theorie de l'information*, Publ. Inst. Statist. Univ. Paris **3**, 3 (1954).
- [24] J. Harvda and F. Charvat, *Kybernetika* **3**, 30 (1967).
- [25] I. Vajda, *Kybernetika* **4**, 105 (1968) [in Czech].
- [26] Z. Daroczy, *Inf. and Control* **16**, 36 (1970).
- [27] B.D. Sharma and D.P. Mittal, *J. Math. Sci.* **10**, 28 (1975).
- [28] A. Wehrl, *Rev. Mod. Phys.* **50**, 221 (1978). See also I.J. Taneja, *Advances in Electronics and Electron Physics* **76**, 327 (1989); M. Behara, *Additive and nonadditive measures of entropy* (Wiley Eastern, New Delhi, 1990); M. Bas-seville, Institut de Recherche en Informatique et Systemes Aleatoires -IRISA (France), Report 1020 (May 1996).
- [29] E.M.F. Curado and C. Tsallis, *J. Phys. A* **24**, L69 (1991) [E: **24**, 3187 (1991) and **25**, 1019 (1992)]; C. Tsallis, R.S. Mendes and A.R. Plastino, *Physica A* **261**, 534 (1998).
- [30] A.M. Mariz, *Phys. Lett. A* **165**, 409 (1992); see also J.D. Ramshaw, *Phys. Lett. A* **175**, 169 (1993).
- [31] A. Chame and E.V.L. de Mello, *J. Phys. A* **27**, 3663 (1994).
- [32] A.K. Rajagopal, *Phys. Rev. Lett.* **76**, 3469 (1996); E.K. Lenzi, L.C. Malacarne and R.S. Mendes, *Phys. Rev. Lett.* **80**, 218 (1998); A.K. Rajagopal, R.S. Mendes and E.K. Lenzi, *Phys. Rev. Lett.* **80**, 3907 (1998).
- [33] F. Buyukkilic and D. Demirhan, *Phys. Lett. A* **181**, 24 (1993); F. Buyukkilic, D. Demirhan and A. Gulec, *Phys. Lett. A* **197**, 209 (1995); U. Tirnakli, F. Buyukkilic and D. Demirhan, *Phys. Lett. A* **245**, 62 (1998); U. Tirnakli and D.F. Torres, *Eur. Phys. J. B* **14**, 691 (2000).
- [34] A.K. Rajagopal and S. Abe, *Phys. Lett. A* **272**, 341 (2000).
- [35] C. Tsallis, S.V.F. Levy, A.M.C. de Souza and R. Maynard, *Phys. Rev. Lett.* **75**, 3589 (1995) [E: **77**, 5442 (1996)]; D. Prato and C. Tsallis, *Phys. Rev. E* **60**, 2398 (1999).
- [36] P.A. Alemany and D.H. Zanette, *Phys. Rev. E* **49**, R956 (1994); D.H. Zanette and P.A. Alemany, *Phys. Rev. Lett.* **75**, 366 (1995); M.O. Caceres and C.E. Budde, *Phys. Rev. Lett.* **77**, 2589 (1996); D.H. Zanette and P.A. Alemany, *Phys. Rev. Lett.* **77**, 2590 (1996).
- [37] A.R. Plastino and A. Plastino, *Physica A* **222**, 347 (1995); C. Tsallis and D.J. Bukman, *Phys. Rev. E* **54**, R2197 (1996); A. Compte and D. Jou, *J. Phys. A* **29**, 4321 (1996); D.A. Stariolo, *Phys. Rev. E* **55**, 4806 (1997).
- [38] L. Borland, *Phys. Lett. A* **245**, 67 (1998); L. Borland, *Phys. Rev. E* **57**, 6634 (1998); L. Borland, F. Pennini, A.R. Plastino and A. Plastino, *Eur. Phys. J. B* **12**, 285 (1999); G. Drazer, H.S. Wio and C. Tsallis, *Phys. Rev. E* **61**, 1417 (2000); A.R. Plastino, M. Casas and A. Plastino, *Physica A* **280**, 289 (2000); M. Bologna, C. Tsallis and P. Grigolini, *Phys. Rev. E* **62**, 2213 (2000).
- [39] T.D. Frank, A. Daffertshofer, P.J. Beek and H. Haken, *Physica D* **127**, 233 (1999).
- [40] F.Y. Edgeworth, *Phil. Mag. (Fifth Series)* **16**, 300 (1883).
- [41] A.M.C. de Souza and C. Tsallis, *Physica A* **236**, 52 (1997).


- [42] S.M. Stigler, *Statistics on the Table - The History of Statistical Concepts and Methods* (Harvard University Press, Cambridge, 1999), p. 284.
- [43] C. Tsallis, A.R. Plastino and W.-M. Zheng, *Chaos, Solitons and Fractals* **8**, 885 (1997); M.L. Lyra and C. Tsallis, *Phys. Rev. Lett.* **80**, 53 (1998); V. Latora, M. Baranger, A. Rapisarda and C. Tsallis, *Phys. Lett. A* (2000), in press [cond-mat/9907412].
- [44] P. Grassberger and M. Scheunert, *J. Stat. Phys.* **26**, 697 (1981); T. Schneider, A. Politi and D. Wurtz, *Z. Phys. B* **66**, 469 (1987); G. Anania and A. Politi, *Europhys. Lett.* **7**, 119 (1988); H. Hata, T. Horita and H. Mori, *Progr. Theor. Phys.* **82**, 897 (1989).
- [45] U. Tirnakli, G.F.J. Ananos and C. Tsallis, preprint (2000) [cond-mat/0005210].
- [46] M. Buiatti, P. Grigolini and L. Palatella, *Physica A* **268**, 214 (1999); J. Yang and P. Grigolini, *Phys. Lett. A* **263**, 323 (1999); S. Montangero, L. Fronzoni and P. Grigolini, preprint (1999) [cond-mat/9911412].
- [47] J.R. Dorfman, *An Introduction to Chaos in Nonequilibrium Statistical Mechanics*, Cambridge Lecture Notes in Physics **14**, eds. P. Goddard and J. Yeomans (Cambridge University Press, Cambridge, 1999).
- [48] F.A. Tamarit, S.A. Cannas and C. Tsallis, *Eur. Phys. J. B* **1**, 545 (1998).
- [49] P. Bak and K. Sneppen, *Phys. Rev. Lett.* **71**, 4083 (1993).
- [50] A.R.R. Papa and C. Tsallis, *Phys. Rev. E* **57**, 3923 (1998).
- [51] P. Jund, S.G. Kim and C. Tsallis, *Phys. Rev. B* **52**, 50 (1995); J.R. Grigera, *Phys. Lett. A* **217**, 47 (1996); S.A. Cannas and F.A. Tamarit, *Phys. Rev. B* **54**, R12661 (1996); L.C. Sampaio, M.P. de Albuquerque and F.S. de Menezes, *Phys. Rev. B* **55**, 5611 (1997); S. Curilef and C. Tsallis, *Phys. Lett. A* **264**, 270 (1999); H.H.A. Rego, L.S. Lucena, L.R. da Silva and C. Tsallis, *Physica A* **266**, 42 (1999); F.A. Tamarit and C. Anteneodo, *Phys. Rev. Lett.* **84**, 208 (2000).
- [52] V. Latora, A. Rapisarda and S. Ruffo, *Phys. Rev. Lett.* **83**, 2104 (1999).
- [53] C. Anteneodo and C. Tsallis, *Phys. Rev. Lett.* **80**, 5313 (1998).
- [54] A. Campa, A. Giansanti, D. Moroni and C. Tsallis, *Long-range interacting classical systems: Universality in mixing weakening*, preprint (2000) [cond-mat/0007104].
- [55] V. Latora and A. Rapisarda, *Dynamical quasi-stationary states in a system with long-range forces*, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Fractals and Solitons* (2001) [cond-mat/0006112].
- [56] M. Antoni and A. Torcini, *Phys. Rev. E* **57**, R6233 (1998).
- [57] X.-P. Huang and C.F. Driscoll, *Phys. Rev. Lett.* **72**, 2187 (1994).
- [58] B.M. Boghosian, *Phys. Rev. E* **53**, 4754 (1996).
- [59] C. Anteneodo and C. Tsallis, *J. Mol. Liq.* **71**, 255 (1997).
- [60] H. Brands, P.H. Chavanis, R. Pasmanter and J. Sommeria, *Phys. Fluids* **11**, 3465 (1999).
- [61] F.M. Ramos, R.R. Rosa and C. Rodrigues Neto, preprint (1999) [cond-mat/9907348].
- [62] T. Arimitsu and N. Arimitsu, *Phys. Rev. E* **61**, 3237 (2000); *J. Phys. A* **33**, L235 (2000); to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Fractals and Solitons* (2001).
- [63] C. Beck, *Physica A* **277**, 115 (2000).
- [64] I. Bediaga, E.M.F. Curado and J. Miranda, *Physica A* **286**, 156 (2000); see also C. Beck, *Physica A* **286**, 164 (2000).
- [65] R. Hagedorn, *Suppl. Nuovo Cimento* **3**, 47 (1965).
- [66] G. Wilk and Z. Wlodarczyk, *Nucl. Phys. B (Proc. Suppl.)* **75A**, 191 (1999); *Phys. Rev. Lett.* **84**, 2770 (2000); to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Fractals and Solitons* (2001) [hep-ph/0004250].
- [67] D.B. Walton and J. Rafelski, *Phys. Rev. Lett.* **84**, 31 (2000).
- [68] D.B. Ion and M.L.D. Ion, *Phys. Rev. Lett.* **81**, 5714 (1998); *Phys. Rev. Lett.* **83**, 463 (1999); *Phys. Rev. E* **60**, 5261 (1999); *Phys. Lett. B* **482**, 57 (2000); to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Fractals and Solitons* (2001).
- [69] G. Kaniadakis, A. Lavagno and P. Quarati, *Phys. Lett. B* **369**, 308 (1996); P. Quarati, A. Carbone, G. Gervino, G. Kaniadakis, A. Lavagno and E. Miraldi, *Nucl. Phys. A* **621**, 345c (1997); G. Kaniadakis, A. Lavagno and P. Quarati, *Astrophysics and space science* **258**, 145 (1998); G. Kaniadakis, A. Lavagno, M. Lissia and P. Quarati, *Proc. 5th International Workshop on Relativistic Aspects of Nuclear Physics* (Rio de Janeiro-Brazil, 1997); eds. T. Kodama, C.E. Aguiar, S.B. Duarte, Y. Hama, G. Odyniec and H. Strobele (World Scientific, Singapore, 1998), p. 193; M. Coraddu, G. Kaniadakis, A. Lavagno, M. Lissia, G. Mezzorani and P. Quarti, in *Nonextensive Statistical Mechanics and Thermodynamics*, eds. S.R.A. Salinas and C. Tsallis, *Braz. J. Phys.* **29**, 153 (1999); A. Lavagno and P. Quarati, *Nucl. Phys. B, Proc. Suppl.* **87**, 209 (2000) (communicated at TAUP99, September 6-10, 1999, Paris); A. Lavagno and P. Quarati, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Fractals and Solitons* (2001).
- [70] A. Lavagno, G. Kaniadakis, M. Rego-Monteiro, P. Quarati and C. Tsallis, *Astrophysical Letters and Communications* **35**, 449 (1998).
- [71] A. Upadhyaya, J.P. Rieu, J.A. Glazier and Y. Sawada, *Anomalous diffusion and non-Gaussian velocity distribution of single Hydra cells in cellular aggregates*, preprint (2000).
- [72] C. Tsallis, F.C. Sa Barreto and E.D. Loh, *Phys. Rev. E* **52**, 1447 (1995).
- [73] C. Tsallis, G. Bemsiki and R.S. Mendes, *Phys. Lett. A* **257**, 93 (1999).
- [74] S. Denisov, *Phys. Lett. A* **235**, 447 (1997); C. Tsallis and M.P. de Albuquerque, *Eur. Phys. J. B* **13**, 777 (2000).
- [75] O. Sotolongo-Costa, A.H. Rodriguez and G.J. Rodgers,


- preprint (2000) [cond-mat/0002339]; preprint (2000) [cond-mat/0004245].
- [76] K.K. Gudima, A.S. Parvan, M. Ploszajczak and V.D. Toneev, preprint (2000) [nucl-th/0003025].
- [77] C. Tsallis and D.A. Stariolo, Notas de Fisica/CBPF 026 (June 1994) and Physica A 233, 395 (1996); D.A. Stariolo and C. Tsallis, Ann. Rev. Comp. Phys., vol. II, ed. D. Stauffer (World Scientific, Singapore, 1995), p. 343.
- [78] T.J.P. Penna, Phys. Rev. E **51**, R1 (1995); K.C. Mundim and C. Tsallis, Int. J. Quantum Chem. **58**, 373 (1996); I. Andricioaei and J.E. Straub, Phys. Rev. E **53**, R3055 (1996); Physica A **247**, 553 (1997); J. Chem. Phys. 107, 9117 (1997); J. Schulte, Phys Rev E **53**, R1348 (1996); J. Schulte, J. Ushio, T. Maruizumi, Thin Solid Films **369**, 285 (2000); K.C. Mundim, T. Lemaire and A. Bassrei, Physica A **252**, 405 (1998); M.A. Moret, P.G. Pascutti, P.M. Bisch and K.C. Mundim, J. Comp. Chemistry **19**, 647 (1998); D.E. Ellis, K.C. Mundim, V.P. Dravid and J.W. Rylander, in *Computer aided-design of high-temperature materials*, 350 (Oxford University Press, Oxford, 1999); D.E. Ellis, K.C. Mundim, D. Fuks, S. Dorfman and A. Berner, Philosophical Magazine B, **79**, 1615 (1999); L. Guo, D.E. Ellis and K.C. Mundim, Journal of Porphyrins and Phthalocyanines **3**, 196 (1999); M.A. Moret, P.M. Bisch and F.M.C. Vieira, Phys. Rev. E **57**, R2535 (1998); A. Berner, K.C. Mundim, D.E. Ellis, S. Dorfman, D. Fuks and R. Evenhaim, Sensors and Actuators **74**, 86 (1999); A. Berner, D. Fuks, D.E. Ellis, K.C. Mundim and S. Dorfman, Applied Surface Science **144-145**, 677 (1999); P. Serra, A.F. Stanton and S. Kais, Phys. Rev. E **55**, 1162 (1997); P. Serra, A.F. Stanton, S. Kais and R.E. Bleil, J. Chem. Phys. **106**, 7170 (1997); Y. Xiang, D.Y. Sun, W. Fan and X.G. Gong, Phys. Lett. A **233**, 216 (1997); U.H.E. Hansmann, Physica A **242**, 250 (1997); U.H.E. Hansmann and Y. Okamoto, Phys. Rev. E **56**, 2228 (1997); U.H.E. Hansmann, Proc. Natl. Acad. Sci. USA **94**, 10652 (1997); U.H.E. Hansmann, Chem. Phys. Lett. **281**, 140 (1997); M.R. Lemes, C.R. Zacharias and A. Dal Pino Jr., Phys. Rev. B **56**, 9279 (1997); P. Serra and S. Kais, Chem. Phys. Lett. **275**, 211 (1997); B.J. Berne and J.E. Straub, Current Opinion in Structural Biology **7**, 181 (1997); H. Nishimori and J. Inoue, J. Phys. A **31**, 5661 (1998); A. Linhares and J.R.A. Torrealo, Int. J. Mod. Phys. C **9**, 133 (1998); R.F. Gutierrez, M. A. de Menezes, C.E. Fellows and O. Dulieu, Chem. Phys. Lett. **300**, 131 (1999); U.H.E. Hansmann, F. Eisenmenger and Y. Okamoto, Chem. Phys. Lett. **297**, 374 (1998); J. E. Straub and I. Andricioaei, in *Algorithms for Macromolecular Modeling*, eds. P. Deuffhard, J. Hermans, B. Leimkuhler, A. Mark, S. Reich and R. D. Skeel, *Lecture Notes in Computational Science and Engineering* **4**, 189 (Springer Verlag, Berlin, 1998); T. Kadowaki and H. Nishimori, Phys. Rev. E **58**, 5355 (1998); J.P.K. Doye, J. Chem. Phys. **109**, 8143 (1998); U.H.E. Hansmann, Y. Okamoto and J.N. Onuchic, Proteins **34**, 472 (1999); U.H.E. Hansmann and Y. Okamoto, Curr. Opin. Struc. Biol. **9**, 177 (1999); Y. Pak and S.M. Wang, J. Chem. Phys. **111**, 4359 (1999); G. Gielis and C. Maes, J. Phys. A **32**, 5389 (1999); M. Iwamatsu and Y. Okabe, Physica A **278**, 414 (2000); Y. Xiang, D.Y. Sun and X.G. Gong, J. Phys. Chem. A **104**, 2746 (2000); Y. Xiang and X.G. Gong, Phys. Rev. E **62** (2000), in press; F. Calvo and F. Spiegelmann, J. Chem. Phys. **112**, 2888 (2000).
- [79] A. Robledo, Phys. Rev. Lett. **83**, 2289 (1999); J. Stat. Phys. **100**, 475 (2000); Granular Matter (2000), in press; A. Robledo and J. Quintana, to appear in the Proc. of the "International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics" (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, Chaos, Fractals and Solitons (2001).
- [80] F.A. Oliveira, B.A. Mello and I.M. Xavier Jr., Phys. Rev. E **61**, 7200 (2000); R.S. Mendes and C. Tsallis, preprint (2000) [cond-mat/0003365].
- [81] G.M. Zaslavsky, Physics Today **52**, 39 (August 1999).
- [82] S. Abe and A.K. Rajagopal, *Nonadditive conditional entropy and its significance for local realism*, Physica A (2000), in press; see also C. Tsallis, S. Lloyd and M. Baranger, preprint (2000) [quant-ph/0007112].
- [83] C. Brukner and A. Zeilinger, *Conceptual inadequacy of the Shannon information in quantum measurements*, quant-ph/0006087.















with $\lim_{N \rightarrow \infty} \tau(N) = \infty$ and $N^* \equiv \frac{N^{1-\alpha/d} - 1}{1 - \alpha/d}$

Fig. 2

